



MOSTAR® 10.8 EC

FACT SHEET

Product Type: Herbicide

Active ingredient: Quizalofop-p-ethyl

Concentration and Formulation: 10.8 % w/v Concentrated Emulsion

Mode of action: Systemic

MAG Registration Number: 4906

Available sizes: 200 cc, 1 L and 20 L.

Main characteristics:

MOSTAR 10.8 EC, is a post-emergent herbicide that acts on annual and perennial grasses, with systemic and selective action for broadleaf crops. The first symptom that appears after foliar application is on growing leaves that yellow and retard growth. It is absorbed through the surface of the leaves and translocated by the plant (xylem-floema), accumulating in the connective tissue.

Crops to be treated:

Pineapple (*Ananas comosus*), Soy (*Glycine max*)

Dosage to be used per crop:

CROP	DOSE	RECOMMENDATIONS
Pineapple (<i>Ananas Commosus</i>)	1.5-2.5 L/ha in 400 L of water	It is recommended to use a volume of 200 to 400 L of water per hectare and a pressure of 60 to 80 pounds per square inch.
Soy (<i>Glycine max</i>)	0.4-0.5 L/ha in 400 L of water	It is recommended to use a volume of 200 to 400 L of water per hectare and a pressure of 60 to 80 lbs/sq. inch.



MOSTAR ® 10.8 EC

FACT SHEET

Period of re-entry to the treated area:

1 day (24 hours).

Application interval:

No more than three applications are recommended during the cycle or handled in accordance with the weed infestation.

Interval between last application and harvest:

15 days.

Compatibility:

It is compatible with most pesticides for agricultural use, except copper-based products.

Precautions:

Care should be taken when applying this product to avoid damaging desirable plants and crops. Do not allow spraying to be carried by wind as it may cause damage to nearby crops, plants and other areas that are not intended to be treated. For handling, use full protective equipment: protective gloves, coveralls, boots, face shield. Do not eat, smoke or drink during handling.