



FACT SHEET

Product Type: Herbicide

Active ingredient: Picloram + 2,4 D

Concentration and Formulation: 30.4 % w/v Liquid Solution

Mode of action: Systemic

MAG Registration Number: 3013

Available sizes: 1 L, 3.5 L, 9.5 L, 19 L and 200 L

Main characteristics:

NAVAJO 30.4 SL is a systemic herbicide based on picloram and 2,4-D, which acts by blocking the normal growth process, which leads to the loss of root function, also produces phloem blocking, photosynthesis inhibition and finally death.

Weeds to be controlled:

Hawthorn (*Acacia farneciana*), Cornishman (*Acacia cornigera*), Mallow (*Anoda cristata*), Viperana (*Asclepias curassavica*), Deer's foot (*Bahuinia excisa*), Charcoal (*Calliandra* spp.), Platanillo (*Canna indica*), Coffee (*Casearia javitensis*), Bean (*Cassia tora*), Brush (*Sida* spp.), Saxophilus (*Sida* spp.), Chipilin de Monte (*Crotalaria retusa*), Golondrina, Lechilla (*Euphorbia* spp.), Guacimo (*Guazuma ulmifolia*), Bijahua (*Helio-cohia bihai*), Guanacaste blanco (*Pitecolobium* spp.), Chile de Perro (*Polygonum* sp.), Chipilin de Monte (*Crotalaria retusa*), Chile de Perro (*Polygonum* sp.)

Crops to be protected:

Pastures.

Dose:

In small weeds of 3-4 true leaves use a dose of 2-4 L/ha. For adult weeds it is necessary to moisten the foliage of the plant well and use a dose of 3-6 L/ha. For control of logs and shrubs use 300 cc - 400 cc of **NAVAJO 30.4 SL** dissolved in 10 liters of water and apply with back pump or brush directly on fresh cuts. The dose per hectare is 3-4 L/ha.



FACT SHEET

General recommendations:

It is recommended to apply it when weeds have 3 to 4 true leaves.

Interval between last application and harvest:

Not applicable as it is applied in the early stages of cultivation.

Interval between last application and grazing:

The safety time between the last application and grazing is 14 days without grazing the cattle after application.

Re-entry interval to the treated area:

You can re-enter the treated area after 24 hours. Otherwise, use the complete protective equipment.

Phytotoxicity:

It is not phytotoxic to crops and in the recommended dose. Do not apply a higher dose than recommended for possible damage to the crop. Care should be taken when applying the product as drifting may damage nearby crops.

Compatibility:

It is compatible with herbicides such as paraquat, diuron, propanil, atrazine and ametrine.

Precautions:

Care should be taken when applying this product to avoid damaging desirable plants and crops. Do not allow spraying to be carried by wind as it may cause damage to nearby crops, plants and other areas that are not intended to be treated. For handling, use full protective equipment: protective gloves, coveralls, boots, face shield. Do not eat, smoke or drink during handling.