



Seraxone

FACT SHEET

Product Type: Herbicide

Active ingredient: Paraquat

Mode of action: Contact

MAG Registration Number: 3013

Available sizes: 1 L, 3.5 L, 19 L and 200 L

Main characteristics:

SERAXONE 20 SL is a post-emergence contact herbicide that acts on all green plant tissues and is particularly active against annual grasses and broadleaf weeds. **SERAXONE 20 SL**, needs active plants to manifest its action, which is characterized by cell wall collapse and desiccation.

Crops to be treated:

Banana, Banana (Musa spp), Coffee (Coffea arabica), Citrus (Citrus spp)

Use in pre-sowing and pre-emergent to the crop:

Potato (Solanum tuberosum), Cabbage (Brassica oleracea var capitata), Carrot (Daucus carota), Onion (Allium cepa), Strawberry (Fragaria spp), Bean (Phaseolus vulgaris), Lettuce (Lactuca sativa), Melon (Cucumis melo), Pineapple (Ananas comosus), Sorghum (Sorghum vulgare), Sugar Cane (Saccharum officinarum), Rice (Oryza sativa), Cotton (Gossypium hirsutum), Corn (Zea mays), Soya (Glycine max)

Dosage to be used per crop:

The dose ranges from 1.5 to 3 liters/ha (90 to 120 ml per 16 liter knapsack sprayer). It is recommended to apply it to weeds that are small, about 15 cm high and actively growing. Do not allow it to spread (very volatile) to neighboring crops, use a protective hood to direct the application only to the weed, especially when it is

between rows.

In the pineapple crop it is recommended to use a dose of 12 to 15 liters/ha for drying. Apply in mixture in a volume of up to 2000 liters of water per hectare.



Seraxone

FACT SHEET

Period of re-entry to the treated area:

When the application has dried on the foliage.

Application interval:

According to the incidence of weeds in the crop.

Interval between last application and harvest:

0 days.

Compatibility:

It is compatible with most pesticides for agricultural use.

Precautions:

Care should be taken when applying this product to avoid damaging desirable plants and crops. Do not allow spraying to be carried by wind as it may cause damage to nearby crops, plants and other areas that are not intended to be treated. For handling, use full protective equipment: protective gloves, coveralls, boots, face shield. Do not eat, smoke or drink during handling.