



HERBICIDA PARA CAÑAS Y POTREROS

FACT SHEET

Product Type: Herbicide

Active ingredient: Picloram + 2,4 D

Concentration and Formulation: 7.5 % w/v Liquid Solution

Mode of action: Systemic

MAG Registration Number: 8702335

Available sizes: 100cc, 250 cc, 1 L, 3.5 L, 19 L and 200 L

Main characteristics:

The product is a systemic herbicide based on picloram and 2,4-D, which acts blocking the normal growth process, which leads to the loss of root function, also produces phloem blocking, photosynthesis inhibition and finally death.

Weeds to be controlled:

Sleepworm (*Mimosa pudica*), Bean (*Cassia tora*), Bledo (*Amaranthus spp.*), Scorpiontail (*Heliotropium indicum*), Verbena (*Stachytarpheta cayennensis*).

Crops to be protected:

Turf and pasture.

GENERAL APPLICATION TO FOLIAGE:

Recommended dose:

4-8 liters/ha. in 400 liters of water

General recommendations:

It is recommended to apply it in areas that present a not very severe infestation of weeds.

PATCHING:

Recommended dose:

1-2 liters/ha. in 200 liters of water.

General recommendations:

It is recommended to apply it directly on the foliage of the weed and when the weed does not cover more than 50% of the pasture.



HERBICIDA PARA CAÑAS Y POTREROS

FICHA TÉCNICA

Interval between last application and harvest:

Not applicable as it is applied in the early stages of cultivation.

Interval between last application and grazing:

You must leave the area without grazing for 14 days after application.

Re-entry interval to the treated area:

You can re-enter the treated area after 24 hours. Otherwise, use the complete protective equipment.

Phytotoxicity:

It is not phytotoxic to crops and at recommended doses. Do not apply a higher dose than recommended for possible damage to the crop. Care should be taken when applying the product as drifting may damage nearby crops.

Compatibility:

It is compatible with herbicides such as paraquat, diuron, propanil, atrazine and ametrine.

Precautions:

Care should be taken when applying this product to avoid damaging desirable plants and crops. Do not allow spraying to be carried by wind as it may cause damage to nearby crops, plants and other areas that are not intended to be treated. For handling, use full protective equipment: protective gloves, coveralls, boots, face shield. Do not eat, smoke or drink during handling.